



Research Article



Fig Juice Fermented with Lactic Acid Bacteria as a Nutraceutical Product

Sima Khezri¹, Parvin Dehghan^{2*}, Razzagh Mahmoudi³, Mahdi jafarlou⁴

¹Department of Food Science and Technology, Faculty of Nutrition and Food Sciences, Student Research Committee, Talented Student Center, Tabriz University of Medical Sciences, Tabriz, Iran.

²Department of Food Science and Technology, Faculty of Nutrition and Food Sciences, Nutrition Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

³Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

⁴Department of Human Anatomy, Faculty of Medicine and Health Sciences, University Putra Malaysia, Serdang, Selangor Darul Ehsan, Malaysia.

Article Info

Article History:

Received: 11 June 2016

Accepted: 14 August 2016

ePublished: 30 December 2016

Keywords:

-Fig juice
-Probiotic
-Lactobacillus
-Nutraceutical
-Survival

ABSTRACT

Background: Probiotics are live microorganisms bringing useful effects to the host through balancing intestine microbiota. This research was undertaken to determine the suitability of fig juice as raw material for production of probiotic juice by three species of lactic acid bacteria (*Lactobacillus casei*, *Lactobacillus plantarum* and *Lactobacillus delbrueckii*).

Methods: Heat treated fig juices were inoculated (6 log CFU/ml) by three species inocula separately and incubated at 30 °C for 72 h. Changes in the pH, acidity, reducing sugar content and viable cell counts during the fermentation were monitored. Sensory characteristics of probiotic fig juice were also evaluated.

Results: *L. delbrueckii* grew well on fig juice; reached nearly 9 log CFU/ml after 48 h of fermentation at 30 °C. After 4 weeks of cold storage at 4 °C, the viable cell counts of *L. delbrueckii* and *L. plantarum* were still 6 and 5 log CFU/ml, respectively, in fermented fig juice; but *L. casei* was just survived until 2th week of cold storage time, reduced from 9 to 3 log CFU/ml. The results of the sensory evaluation showed that fermented fig juice samples were significantly different ($P<0.05$) from the control sample in taste, odor, consistency and overall acceptability. *L. casei* was more acceptable comparing to the others.

Conclusion: *L. delbrueckii* was the most suitable strain from the point of survivability among other species at the consumption time. Therefore, probiotic fig juice can serve as healthy beverage for vegetarians and consumers with lactose-allergy.